

- 2 Coronavirus COVID-19 Global Cases by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU). <https://gisanddata.maps.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b48e9ecf6> (accessed March 19, 2020).
- 3 The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. The epidemiological characteristics of an outbreak of 2019 novel coronavirus disease (COVID-19)—China. 2020. *China CDC Wkly* 2020; **2**: 113–22.
- 4 Negin J, Abimbola S, Marais BJ. Tuberculosis among older adults—time to take notice. *Int J Infect Dis* 2015; **32**: 135–37.
- 5 Wingfield T, Tovar MA, Datta S, Saunders MJ, Evans CA. Addressing social determinants to end tuberculosis. *Lancet* 2018; **391**: 1129–32.
- 6 Saleem A, Azher M. The next Pandemic - Tuberculosis: The oldest disease of mankind rising one more time. *Br J Med Pract* 2013; **6**: a615.
- 7 Gilbert M, Pullano G, Pinotti F, et al. Preparedness and vulnerability of African countries against importations of COVID-19: a modelling study. *Lancet Infect Dis* 2020; **395**: 871–77.
- 8 Meghji J, Lesosky M, Joekes E, et al. Patient outcomes associated with post-tuberculosis lung damage in Malawi: a prospective cohort study. *Thorax* 2020; **75**: 269–78.
- 9 Horton KC, MacPherson P, Houben RMGJ, White RG, Corbett EL. Sex differences in tuberculosis burden and notifications in low- and middle-income countries: a systematic review and meta-analysis. *PLoS Med* 2016; **13**: e1002119.
- 10 Wenham C, Smith J, Morgan R, the Gender and COVID-19 Working Group. COVID-19: the gendered impacts of the outbreak. *Lancet* 2020; **395**: 846–48.



COVID-19: Respiratory support outside the intensive care unit



Al Photo/Science Photo Library

The optimal mode of respiratory support for individuals with severe coronavirus disease 2019 (COVID-19) before invasive mechanical ventilation (IMV) is currently a subject of much debate. Recently published guidelines¹ and a Comment² differ substantially to other guidelines in this regard, with some advocating high flow nasal cannulae (HFNC) over non-invasive ventilation (NIV),^{1,2} or vice versa (NHS guidance). This debate is understandable given the paucity of data and need for rapid generation of guidance, but it is a cause of confusion among respiratory physicians.

Much of the data guiding practice in this area derive from the critical care setting. In acute respiratory distress syndrome (ARDS), early intubation was associated with survival benefit when PaO₂/FiO₂ ratio was <150 when compared with NIV.³ Some COVID-19 guidelines have therefore suggested NIV as a bridging therapy only, before transfer to the intensive care unit (ICU) and IMV. However, up to 50% of patients with COVID-19 admitted to the ICU did not subsequently require IMV.⁴ Given the considerable resource limitations imposed by the current unprecedented viral pandemic, it is important to ascertain whether selected patients can be safely managed outside of the ICU.

To our knowledge, there have been no randomised control trials in the use of either HFNC or NIV in coronavirus-related pneumonia. It has been reported that use of NIV during the Middle Eastern respiratory syndrome outbreak was associated with a 92% risk of requiring IMV, thus suggesting futility.⁵ This study was based in the ICU, however, and patients enrolled had a median PaO₂/FiO₂ ratio of 110 (IQR 62–160), indicating a degree of severity that likely warranted initial

management with IMV. Conversely, data from only one study⁶ on the severe acute respiratory syndrome (SARS) outbreak suggest that NIV can successfully avoid intubation.

HFNC has received much interest since the FLORALI trial.⁷ Acute hypoxaemic respiratory failure (AHRF) in this study was largely secondary to community or hospital-acquired pneumonia. Though the primary outcome of intubation at day 28 was negative, HFNC reduced requirement for intubation in a subgroup of patients with PaO₂/FiO₂ ratio <200 and was associated with a reduction in mortality when compared with NIV or regular oxygen face mask. The NIV group of this study involved NIV use for an average of only 8 h per day, however, and a relatively high target tidal volume of 7–10 mL/kg. FLORALI also utilised a flow rate of 50 L/min with HFNC. To ameliorate potential aerosol generation, a flow limit of 30 L/min in COVID-19 has been proposed. The level of positive end-expiratory pressure (PEEP) supplied is consequently reduced. Notably, with regards to aerosol generation and risk to health-care workers, intubation poses a greater risk than NIV and a risk with HFNC has not been established.⁸

Concern regarding ward oxygen flow rates and hospital oxygen reserves is probably the most important cause for hesitancy over advocating HFNC (Irish Thoracic Society Guidelines). A major benefit of PEEP is that it might allow for down-titration of FiO₂, mitigating against over-consumption of hospital oxygen supply and avoiding hyperoxia-related lung injury. Anecdotal reports and our own experience of COVID-19-related lung injury suggests a good response to application of PEEP, perhaps related

Published Online

April 9, 2020

[https://doi.org/10.1016/S2213-2600\(20\)30176-4](https://doi.org/10.1016/S2213-2600(20)30176-4)

For **NHS guidelines** see <https://www.england.nhs.uk/coronavirus/secondary-care/other-resources/specialty-guides/#adult-critical-care>

For **Irish Thoracic Society guidelines** see <https://irishthoracicsociety.com/2020/03/irish-thoracic-society-covid-19-guidelines-for-managing-respiratory-care/>

For **ARDSnet guidelines** see <https://www.thoracic.org/statements/resources/cc/ards-guidelines.pdf>

to recruitment of atelectasis and reduced work of breathing. PEEP of 10 cm H₂O or higher can shift the lung to the point on the pressure–volume curve with the highest slope (high compliance).⁹ Haemodynamic instability appears to be a relatively infrequent feature of these patients, and thus higher PEEP (ie, expiratory positive airway pressure) than traditionally applied with NIV is likely to be tolerated well.

Benefits of bi-level positive airway pressure over continuous positive airway pressure in this setting have not been established conclusively. Regardless of mode, the key factor in improving oxygenation is mean airway pressure (Paw). Addition of pressure support has the advantage of compensating for resistance present in the tubing and in further reducing work of breathing.¹⁰ It is prudent to follow ARDSnet guidelines in maintaining tidal volume of ≤6 mL/kg through low pressure support (driving pressure), relatively high PEEP, and the lowest FiO₂ feasible. To mitigate against nosocomial aerosol transmission, it is critical that NIV circuits are modified to include a filter at the exhalation port or vent.

The debate about the optimal mode of respiratory support before IMV in AHRF has not been settled, much less in the setting of coronavirus, and it is important to note that harm can be caused if inappropriate treatment is used.³ Evidence from China¹¹ suggests that a large minority of patients with severe respiratory failure due to SARS coronavirus 2 (SARS-CoV-2) can avoid intubation via use of NIV however. NIV is a well-established therapy with which general respiratory physicians and nurses are familiar, and which is readily applicable in the non-critical care setting. Caveats would

include careful patient selection so as not to delay IMV where appropriate, modified settings specific to the pathophysiology of COVID-19, and mitigation against infection transmission by aerosol.

RWC reports grants and personal fees from Aerogen and GlaxoSmithKline, personal fees from Novartis and TEVA, and grants from Vitalograph, outside of the submitted work. TM and CG declare no competing interests.

*Tom McEnery, Ciara Gough, Richard W Costello
thomasmcenery@rcsi.ie

Royal College of Surgeons in Ireland, Beaumont Hospital, Dublin D09 V2N0, Ireland

- 1 Alhazzani W, Moller MH, Arabi YM, et al. Surviving Sepsis Campaign: guidelines on the management of critically ill adults with Coronavirus Disease 2019 (COVID-19). *Intensive Care Med* 2020; published online March 28. DOI:10.1007/s00134-020-06022-5.
- 2 Matthay MA, Aldrich JM, Gotts JE. Treatment for severe acute respiratory distress syndrome from COVID-19. *Lancet Respir Med* 2020; published online March 20. [https://doi.org/10.1016/S2213-2600\(20\)30127-2](https://doi.org/10.1016/S2213-2600(20)30127-2).
- 3 Bellani G, Laffey JG, Pham T, et al. Non-invasive ventilation of patients with acute respiratory distress syndrome: insights from the LUNG SAFE study. *Am J Respir Crit Care Med* 2017; **195**: 67–77.
- 4 Meng L, Qiu H, Wan L, et al. Intubation and ventilation amid the COVID-19 outbreak: wuhan's experience. *Anesthesiol* 2020; published online March 26. DOI:10.1097/ALN.0000000000003296.
- 5 Alraddadi BM, Qushmaq I, Al-Hameed FM, et al. Non-invasive ventilation in critically ill patients with the Middle East respiratory syndrome. *Influenza Other Respir Viruses* 2019; **13**: 382–90.
- 6 Cheung TM, Yam LY, So LK, et al. Effectiveness of non-invasive positive pressure ventilation in the treatment of acute respiratory failure in severe acute respiratory syndrome. *Chest* 2004; **126**: 845–50.
- 7 Frat JP, Thille AW, Mercat A, et al. High-flow oxygen through nasal cannula in acute hypoxemic respiratory failure. *N Engl J Med* 2015; **372**: 2185–96.
- 8 Fowler RA, Guest CB, Lapinsky SE, et al. Transmission of severe acute respiratory syndrome during intubation and mechanical ventilation. *Am J Respir Crit Care Med* 2004; **169**: 1198–202.
- 9 L'Her E, Deye N, Lellouche F, et al. Physiologic effects of non-invasive ventilation during acute lung injury. *Am J Respir Crit Care Med* 2005; **172**: 1112–18.
- 10 Katz JA, Marks JD. Inspiratory work with and without continuous positive airway pressure in patients with acute respiratory failure. *Anesthesiol* 1985; **63**: 598–607.
- 11 Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020; **323**: 1061–69.

Hydroxychloroquine in the management of critically ill patients with COVID-19: the need for an evidence base



With the rapid spread of the novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), critical care physicians are seeing increasing numbers of patients with acute respiratory failure secondary to coronavirus disease 2019 (COVID-19) and reporting mortality rates of 40–65% for those requiring mechanical ventilation¹—strikingly higher than the mortality rates reported for the more typical acute respiratory distress syndrome associated with other

diseases.² The focus of therapeutic intervention has therefore been not only to reverse hypoxaemia and provide adequate organ support, but also to decrease viral load and thus limit disease severity. In addition to several antiviral agents, antimalarial drugs have been proposed as treatments that could reduce transmission of the virus. In-vitro studies have shown that chloroquine and hydroxychloroquine can both inhibit SARS-CoV-2 transmission,^{3–5} through

Published Online
April 15, 2020
[https://doi.org/10.1016/S2213-2600\(20\)30172-7](https://doi.org/10.1016/S2213-2600(20)30172-7)